



## The 2008 Antarctic Ozone Hole Summary: Monday 8 December 2008

Paul Krummel<sup>1</sup>, Paul Fraser<sup>1</sup> and Paul Lehmann<sup>2</sup>

Centre for Australian Weather and Climate Research

<sup>1</sup>CSIRO Marine & Atmospheric Research, Aspendale, Victoria

<sup>2</sup>Bureau of Meteorology, Melbourne, Victoria

### Instrumentation

Data from the Ozone Monitoring Instrument (OMI) on board the Earth Observing Satellite (EOS) Aura, processed with the NASA TOMS Version 8.5 algorithm, are being utilized again this year in our weekly ozone hole reports. OMI continues the NASA TOMS satellite record for total ozone and other atmospheric parameters related to ozone chemistry and climate.

### The 2008 ozone hole

The data show that the ozone minimum dipped to 100 DU by early October, about a week later than in 2005 (Figure 1, top panel), and has since recovered to 155 DU by the end of the 2<sup>nd</sup> week of November, staying at that value until the end of the 3<sup>rd</sup> week of November and reaching almost 160 DU by the end of November. As expected, a rapid recovery of ozone was seen in the hole during the first week of December, reaching 190 DU by December 5, similar in recovery to the 2006 hole.

The ozone hole reached a maximum area of 27 million km<sup>2</sup> for 2008 in early October, declined to 17 million km<sup>2</sup> by the end of October (Figure 1, bottom panel), increased again to 17-18 million km<sup>2</sup> during the first week of November, declining rapidly to 15-16 million km<sup>2</sup> by the end of the 2<sup>nd</sup> week of November. By the end of the 3<sup>rd</sup> week of November the area increased slightly to 16-17 million km<sup>2</sup>, decreasing again to 14-15 million km<sup>2</sup> by the end of November. The ozone hole area remained quite stable over the 3-4 weeks at 15-18 million km<sup>2</sup>. However, over the past 2 weeks the area of the ozone hole has dropped dramatically to 6 million km<sup>2</sup> by December 5, signalling the near-end of this year's hole.

The ozone deficit grew to a maximum of 40 Mt by early October, declining to 12-13 Mt by the end of the 2<sup>nd</sup> week of November (Figure 2, top panel) and it has stayed at 12-13 Mt by the end of the 3<sup>rd</sup> week of November, falling sharply to 10 Mt by the end on November and even more rapidly to 2 Mt by December 5. This rapid decline in deficit has occurred when the vortex has again become distorted (see below).

The average ozone amount in the hole (averaged column ozone amount in the hole weighted by area, Figure 2 bottom panel) fell to 150 DU by early October, and has since recovered to 180-185 DU by the end of 2<sup>nd</sup> week of November, staying at that value until the end of the 3<sup>rd</sup> week of November and rising to 190 DU by the end of November. The rapid ozone rise continued in the first week of December, reaching about 205 DU by December 5.



At present the 2008 ozone hole is as persistent as any that have gone before. The next few weeks will decide whether it is the most persistent ever. The data suggest a rapid closure from now on (as in 2006) may occur.

Total column ozone data over Australia and Antarctica for 24 November to 5 December are shown in Figure 3. The Antarctic coast south of Australia has been outside the hole for this entire period. The vortex was relatively symmetrical from 16-22 November. This symmetrical nature is what is caused the persistence of this year's hole. The symmetrical nature lasted until 22 November and since then the vortex has become significantly distorted along the 30°E-120°W axis. The South Pole is now on the edge of the hole. Due to this distortion, the ozone hole again passed over the southern tip of South America on 27 November. Over the period December 2-5 the hole became less distorted but centred on the Antarctic coast south of Africa.

It is now possible to rank the 2008 Antarctic ozone hole for all the metrics we calculate except integrated ozone deficit, which we have estimated in the rankings below (this metric can be finalised once the ozone hole has closed).


Of the 29 holes for which we have data since 1979, the 2008 hole ranked

6 <sup>th</sup> :	15-day average area	11 <sup>th</sup> :	15-day average minimum ozone
7 <sup>th</sup> :	daily maximum area	10 <sup>th</sup> :	daily minimum ozone
6 <sup>th</sup> :	daily ozone deficit	8 <sup>th</sup> :	daily min average ozone amount in the hole
~7 <sup>th</sup> :	integrated ozone deficit (estimated)		




Last updated on Mon Dec 08 12:25:46 2008 by kru021@PBKNB-AS

**Figure 1:** Ozone hole depth (top panel) and area (bottom panel) based on OMI satellite data, as of 5 December 2008.



Last updated on Mon Dec 08 12:25:46 2008 by kru021@PBKNB-AS

**Figure 2:** Estimated daily ozone deficit (top panel) and average ozone amount within the ozone hole (bottom panel) based on OMI satellite data, as of 5 December 2008.



**Figure 3:** OMI ozone hole images for 24 November to 5 December 2008; the ozone hole boundary is indicated by the red 220 DU contour line.

## Definitions

CFCs: chlorofluorocarbons, synthetic chemicals containing chlorine, once used as refrigerants, aerosol propellants and foam-blowing agents, that break down in the stratosphere (15-30 km above the earth's surface), releasing reactive chlorine radicals that catalytically destroy stratospheric ozone.

DU: Dobson Unit, a measure of the total ozone amount in a column of the atmosphere, from the earth's surface to the upper atmosphere, 90% of which resides in the stratosphere at 15 to 30 km.

Halons: synthetic chemicals containing bromine, once used as fire-fighting agents, that break down in the stratosphere releasing reactive bromine radicals that catalytically destroy stratospheric ozone. Bromine radicals are about 50 times more effective than chlorine radicals in catalytic ozone destruction.

Ozone: a reactive form of oxygen with the chemical formula  $O_3$ ; ozone absorbs most of the UV radiation from the sun before it can reach the earth's surface.

Ozone Hole: ozone holes are examples of severe ozone loss brought about by the presence of ozone depleting chlorine and bromine radicals, whose levels are enhanced by the presence of PSCs (polar stratospheric clouds), usually within the Antarctic polar vortex. The chlorine and bromine radicals result from the breakdown of CFCs and halons in the stratosphere. Smaller ozone holes have been observed within the weaker Arctic polar vortex.

Polar night terminator: the delimiter between the polar night (continual darkness during winter over the Antarctic) and the encroaching sunlight. By the first week of October the polar night has ended at the South Pole.

Polar vortex: a region of the polar stratosphere isolated from the rest of the stratosphere by high west-east wind jets centred at about  $60^\circ S$  that develop during the polar night. The isolation from the rest of the atmosphere and the absence of solar radiation results in very low temperatures (less than  $-78^\circ C$ ) inside the vortex.

PSCs: polar stratospheric clouds are formed when the temperatures in the stratosphere drop below  $-78^\circ C$ , usually inside the polar vortex. This causes the low levels of water vapour present to freeze, forming ice crystals and usually incorporates nitrate or sulphate anions.

TOMS & OMI: the Total Ozone Mapping Spectrometer & Ozone Monitoring Instrument, are satellite borne instruments that measure the amount of back-scattered solar UV radiation absorbed by ozone in the atmosphere; the amount of UV absorbed is proportional to the amount of ozone present in the atmosphere.

UV radiation: a component of the solar radiation spectrum with wavelengths shorter than those of visible light; most solar UV radiation is absorbed by ozone in the stratosphere; some UV radiation reaches the earth's surface, in particular UV-B which has been implicated in serious health effects for humans and animals; the wavelength range of UV-B is 280-315 nanometres.

## Acknowledgements

The TOMS and OMI data are provided by the TOMS ozone processing team, NASA Goddard Space Flight Center, Atmospheric Chemistry & Dynamics Branch, Code 613.3. The OMI instrument was developed and built by the Netherlands's Agency for Aerospace Programs (NIVR) in collaboration with the Finnish Meteorological Institute (FMI) and NASA. The OMI science team is lead by the Royal Netherlands Meteorological Institute (KNMI) and NASA.